

Cytotoxic T-Lymphocyte Antigen 4 (*CTLA-4*) Gene Variants in Females with Recurrent Spontaneous Pregnancy Loss

Osama Saad Elshaer¹, Eman Gamal Behiry¹, Hesham Ali Issa¹, Dalia Farouk Hussen², Engy A. Ashaat³, Weaam Mohammed Mohammed El-kholey⁴, Mohammed Abd Elmaboud El Gazzar⁵, Abeer Ramadan⁶

Original
Article

¹Clinical and Chemical Pathology Department, ⁵Obstetrics and Gynecology Department, Faculty of Medicine, Benha University, ²Human Cytogenetic Department, ³Clinical Genetics Department, ⁶Molecular Genetics and Enzymology Department, Human Genetics and Genome Research Institute, National Research Centre, ⁴Clinical Pathology Department, Faculty of Medicine, Benha University, Egypt.

ABSTRACT

Background: Merely 2% of pregnant mothers experience two consecutive miscarriages, and as many as 50% of patients suffering from Recurrent Pregnancy Loss (RPL) lack a definitive cause for their condition. The *CTLA-4* gene is located in band q33 on chromosome 2 of the human genome, is approximately 6.2 kilobases long, and is composed of 3 introns and 4 exons. In decidual and peripheral dendritic cells, it is expressed on human placental regulatory T (Treg) cells. Human miscarriages were associated with a downregulation of Treg cells and *CTLA-4* expression in decidual and peripheral lymphocytes. The purpose of our analysis was to determine whether the *CTLA-4* +49A/G (rs231775) & (rs3087243) gene polymorphism and unexplained RPL were related.

Subjects and Methods: This case-control study included women with RPL were contrasted with healthy females at the age of motherhood. The study was conducted at the National Research Centre (Clinical Genetics Department -Human Cytogenetic Department - Molecular Genetics and Enzymology Department) in collaboration with Benha Faculty of Medicine (Clinical and Chemical Pathology Department).

Results: Considering AG is the reference haplotype, no significant association was found between *CTLA-4* rs231775-rs3087243 haplotypes with number of abortions. rs231775 AG+GG was considered a protective predictor, while rs3087243 GA+AA was considered a risky predictor of susceptibility to spontaneous recurrent abortion in uni- and multivariable analyses.

Conclusion: Our study revealed that a significant correlation was discovered between the *CTLA-4* gene rs231775 AG, GG genotypes, G allele, rs231775- rs3087243 GG haplotype with protective effect against RPL. Whereas, a significant correlation was discovered between the *CTLA-4* gene rs3087243 GA, AA genotypes, A allele AA haplotype and RPL risk.

Key Words: *CTLA-4*; Gene Polymorphism; Recurrent spontaneous abortion.

Received: 31 December 2023, **Accepted:** 8 September 2024

Corresponding Author: Weaam Mohammed Mohammed El-kholey, M.B.B.CH., Faculty of Medicine, Benha University, Egypt. **Tel.:** 01147010214,

E-mail: weamslkholy@gmail.com

ISSN: 2090-8571, 2023

INTRODUCTION

Recurrent pregnancy loss (RPL) is characterized as three or more consecutive spontaneously failed clinical pregnancies before 24 weeks of pregnancy, according to the Royal College of Obstetricians and Gynecologists (RCOG) (Pourmasumi *et al.*, 2018).

There are several terms used to describe repeated pregnancy disruptions, including recurrent spontaneous abortion (RSA), recurrent miscarriage (RM), and RPL.

Numerous studies have revealed that roughly one in every 100 women encounter RPL (Turesheva *et al.*, 2023).

2 to 5% of RPL cases are caused by genetic factors, which can include gene mutations or chromosomal abnormalities. An increasing number of research have concentrated on genetic variables, namely single nucleotide polymorphism (SNP) (Abdi-Shayan *et al.*, 2016).

The mother-fetus relationship is important for a successful pregnancy. Since the fetus and the mother do not share the same genetic makeup, this relationship is essential. It has been discovered that this fetomaternal connection is established by multiple pathways. One of the most significant of these is immunological mechanisms, which enable the mother immune system to develop tolerance toward the semiallogeneic fetus (Mumusoglu *et al.*, 2016).

It is known that the tolerance of the fetus is influenced by negative co-stimulatory signalling pathways, such as *CTLA-4/B7*, which are expressed on Tregs and regulate effector T cell activation and proliferation (Nasiri and Rafti, 2016).

These signalling molecules function by inhibiting immune-reactive cells, and it has been demonstrated that reduced *CTLA-4* expression contributes to human foetal loss (Tripathi and Guleria, 2015).

The *CTLA-4* gene comprises four exons and more than 100 polymorphism sites and it corresponds to human chromosome 2q33. Of these, *CTLA-4 +49A/G* (rs231775) was examined in great detail and has been linked to a heightened risk of developing several human diseases, including Gravis diseases, RPL, and Type 2 diabetes mellitus (Rafti and Nasiri, 2016).

The study's objective was to assess the relationship of unexplained RPL with *CTLA-4 +49A/G* (rs231775) & (rs3087243) gene polymorphisms (SNPs).

SUBJECTS AND METHODS:

Subject:

This case-control study included women with RPL were contrasted with healthy females at the age of motherhood. The study was conducted at the National Research Centre (Clinical Genetics Department- Human Cytogenetic Department - Molecular Genetics and Enzymology Department) in collaboration with Benha, Faculty of Medicine (Clinical and Chemical Pathology Department).

Fifty women in the study group, ages 20 to 35, had undergone at least three spontaneous abortions prior to the 24-week gestational period. Fifty women who were of the same age as the study group, had given birth to at least one healthy child, and had no prior history of pregnancy loss comprised the control group.

We excluded any abnormality including US as uterine abnormalities, thrombophilia screening, immunological testing, any endocrine problem including the thyroid and Karyotyping for both couples was performed.

Controls and research subjects were matched with regard to all other conceivable characteristics (There were no users of oral contraceptives, hormonal medications, or any other dangerous drugs that could have impacted vital bodily functions).

Those with established RPL causes as uterine structural abnormalities, Immunological causes, Medical causes as hypertension, Diabetes mellitus and Chromosomal abnormalities are excluded from this study.

Every study sample was drawn from Clinical Genetics Department at the National Research Centre from May 2018 to October 2019 and the control samples were recruited from Obstetric and Gynecology Outpatient Clinic, Benha University Hospital, from May 2018 to October 2019.

The National Research Center's Ethical Medical Committee and Benha University approved the study protocol. Each and every participant in the study gave their informed consent.

Methods:

The following was conducted on participants:

A thorough history and clinical evaluation paying attention to:

Age and history of recurrent pregnancy loss with no uses of oral contraception, hormonal or any serious medications.

Clinical assessment included abnormal and pelvic U/S.

Hystoscopy to detect normal uterine cavity.

Detailed pregnancy and obstetric history, medical history, family history, past history and consanguinity.

Cytogenetic analysis:

G-banding karyotyping for peripheral blood chromosomes was performed for all patients to exclude any patient with chromosomal abnormality from the study Verma and Babu, (VERMA and BABU, 1995). Approximately 30 metaphases were analyzed and karyotyped for each patient following the International System for Human Cytogenomic Nomenclature [ISCN, 2020] recommendations (MCGOWAN-JORDAN *et al.*, 2020).

Molecular investigations:

Sampling:

Five millilitres of venous blood were collected in EDTA-containing tubes (violet topped) under aseptic precautions and storage at -40°C to measure haemoglobin and identify SNPs utilizing Polymerase Chain Reaction-Restriction Fragment Polymorphism (PCR/RFLP).

Analysis of the polymorphism in CTLA4+49A/G (rs231775) and (rs3087243) by PCR-RFLP method with the following steps:

A. Extraction of genomic DNA from EDTA blood leucocytes.

B. Amplification of the isolated by sets of primers which detect the targeted polymorphism.

C. Detection of PCR amplified products using gel electrophoresis on 2% agarose gel with Tris borate EDTA buffer at 100 V and visualisation was done by ultraviolet (UV) light trans-illumination.

D. Digestion of PCR amplification products was done using the suitable restriction enzyme Endonucleas BbNI (Fermentase, Burlington, USA) specific for target polymorphism.

E. The past digestion products were examined by gel electrophoresis on 2% agarose gel comprising ethidium bromide and visualized by trans illumination for genotype calling.

Determination of CTLA-4+49A/G (rs231775) & (rs3087243) Extraction of DNA:

The purification of the genomic DNA of whole blood DNA was extracted from peripheral blood leucocytes using a Mini Kit employing a Wizard genomic DNA purification kit.

DNA amplification:

Enzymatic amplification was performed by PCR using PCR GOtaq polymerase enzyme (Madison, USA) and veriti thermal cycler (AB Applied Biosystem, Singapore).

Reagents:

- Green Master Mix (Madison, USA) contains i-TaqDA polymerase, reaction buffer, d NTPs, d CTP, d GTP, d TTP). MgCl₂.
- CTLA4-49 (rs231775)&(rs3087243).

Protocol for PCR amplification:

For each reaction, the following were added to the Amico-centrifuge tube:

- DNA template.....2 µL
- I-Star Taq DNA polymerase.....0.5 µL
- The exon 1 position 49 polymorphic restriction sites of the CTLA4 gene were genotyped via PCR amplification; the restriction fragment length polymorphism of the *CTLA-4* gene was then amplified via PCR using the subsequent primers: (rs231775) forward TCCTGAAGACCTGAACACCG and (rs231775) reverse TGCCTTTGACTGCTGAAAC.
- (rs3087243) forward AGGCAGCAGGTGGCAGAAT and (rs3087243) reverse TAAGCAAGAATCACAGAGGGC

The following parameters were utilized for the amplification: a 3-minute initial denaturation at 94, followed by 35 cycles of 45-second denaturation at 94.

Detection of the band of amplified material use trans-illuminating UV nation:

For the existence of the amplification band at 88 and 74 bp, the gel was observed through the filter region of an Alpha Innotech corporation transilluminator.

Gel was then photographed:

- *Left lane:* PCR marker (50-1000 bp Ladder).
- *Other lanes:* presence of the amplification band at 88 and 74bp.

Digestion of the amplified products by restriction enzyme:

CTLA-4 (restriction enzyme) (Biolab, NewEngland) used for digestion of the amplified targeted DNA to study rs231775 & rs3087243.

Detection of the CTLA-4 gene polymorphism:

The *CTLA-4* polymorphism was detected using 2% agarose gel electrophoresis.

Statistical methods:

The data that had been gathered was reviewed, coded, and tabulated utilizing the Statistical Package for the Social Sciences (IBM Corp., 2017). Version 25.0 of IBM SPSS Statistics for Windows (Armonk, NY: IBM Corp). Data were shown, and appropriate analysis was carried out based on the kind of data found for each parameter.

RESULTS

Cytogenetic analysis revealed normal outcomes for all patients; 46,XX (Figure 1).

The current molecular investigation involved 50 cases of unexplained recurrent abortions. Their mean (\pm SD) age was 23.8 (\pm 2.6) years. Furthermore, 50 age-matched, healthy control females ($p > 0.05$) (Table 1).

Considering AA and A are the reference genotype and allele respectively, AG, GG, AG+GG genotypes, G allele exhibited a significantly reduced incidence of in contrast to control groups ($p < 0.05$ for each), and had a protective effect against recurrent unexplained abortions (OR<1 for each) (Table 2).

Considering GG is the reference genotype and G is the reference allele, GA, AA genotypes, dominant model and A allele had significantly greater proportions in cases in contrast to control group ($p < 0.05$ for each), with risk to develop recurrent abortions (OR>1 for each) (Table 3).

Haplotype is a group of alleles that are inherited together from a single parent. The rs231775 and rs3087243 SNPs are spaced about 6 kilo base pairs apart. The rs231775-rs3087243 haplotypes were calculated among each group, AG haplotype showed the highest frequency

among cases (38.7%), and controls (45.9%), GG haplotype displayed the least frequency among cases (10.3%), while AA haplotype displayed the least frequency among controls (5.1%). Considering AG is the reference haplotype, GG showed lower frequency in cases when compared to controls ($p=0.015$), with protective effect against recurrent abortion development ($OR<1$), while AA showed greater frequency in cases in contrast to controls ($p<0.001$), with susceptibility risk for recurrent abortion ($OR>1$). GA haplotype was not associated significantly with recurrent abortion (Table 4).

Considering AG is the reference haplotype, no significant association was found between *CTLA-4* rs231775-rs3087243 haplotypes with number of abortions (Table 5).

Regression analysis was done in order to predict spontaneous recurrent abortion susceptibility utilizing rs3087243, rs231775 genotypes as covariates. rs231775 AG+GG was considered a protective predictor, while rs3087243 GA+AA was considered a risky predictor of susceptibility to spontaneous recurrent abortion in uni- and multivariable analyses (Table 6).

When alleles at two or more loci in a population are not randomly associated, it is referred to as LD. The value of D' ranges from 0 (no disequilibrium) to 1. (maximum disequilibrium). Among our studied control group, $D'=0.91$; while among our studied cases, $D'=0.31$. Our results in control group agreed with those who demonstrated a strong linkage disequilibrium between rs231775, and rs3087243 (Figure 2).

When alleles at two or more loci in a population are not randomly associated, it is referred to as linkage disequilibrium (LD). The value of D' ranges from 0 (no disequilibrium) to 1. (maximum disequilibrium). Among studied control group, $D'=0.91$; while among studied cases, $D'=0.31$ (Figure 2).

Table 1: Comparison of age between studied groups:

	Control N=50	Cases N=50	P
	Mean±SD	Mean±SD	
Age (years)	24.8±2.7	23.8±2.6	0.881

SD: standard deviation; Numerical data are expressed as mean±SD, compared by *t* test.

Table 2: Association of CTLA-4 rs231775 genotypes and alleles with studied groups:

		Control N=50	Cases N=50	p	OR (95% CI)
		N (%)	N (%)		
rs231775	AA	10(20%)	21(42%)	-	Reference
	AG	31(62%)	24(48%)	0.047*	0.438(0.194-0.988)
	GG	9(18%)	5(10%)	0.032*	0.538(0.305-0.947)
	AG+GG	40(80%)	29(58%)	0.018*	0.516(0.299-0.892)
	A	51(51%)	66(66%)	-	Reference
	G	49(49%)	34(34%)	0.031*	0.677(0.475-0.966)

Data is presented as frequency (percentage); *: Significant as p -value <0.05 ; OR: odds ratio; CI: confidence interval. Logistic regression analysis was used. AA is the reference genotype and A is the reference allele; AG+GG is the dominant model.

Table 3: Association of CTLA-4 rs3087243 genotypes and alleles with studied groups:

		Control N=50	Cases N=50	p	OR (95% CI)
		N (%)	N (%)		
rs3087243	GG	21(42%)	9(18%)		Reference
	GA	27(54%)	31(62%)	0.036*	1.842(1.04-3.263)
	AA	2(4%)	10(20%)	0.002*	4.445(1.691-11.688)
	GA+AA	29(58%)	41(82%)	0.009*	2.098(1.202-3.661)
	G	69(69%)	49(49%)		Reference
	A	31(31%)	51(51%)	0.004*	1.69(1.181-2.417)

Data is presented as frequency (percentage); *: Significant as p -value <0.05 ; OR: odds ratio; CI: confidence interval. Logistic regression analysis was used. GG is the reference genotype and G is the reference allele; GA+AA is the dominant model.

Table 4: Association of CTLA-4 rs231775-rs3087243 haplotypes among studied groups:

rs231775-rs3087243	Control	Cases	<i>p</i>	OR (95% CI)
	Frequencies	Frequencies		
AG	0.459	0.387	-	Reference
GA	0.259	0.237	0.721	1.072(0.576-1.995)
GG	0.231	0.103	0.015*	0.656(0.315-0.967)
AA	0.051	0.273	<0.001*	2.854(1.287-6.324)

OR: odds ratio; CI: confidence interval. Logistic regression analysis was used.

Table 5: Association of CTLA-4 rs231775-rs3087243 haplotypes with number of abortions:


rs231775-rs3087243	3 abortions	>3 abortions	<i>p</i>	OR (95% CI)
	Frequencies	Frequencies		
AG	0.404	0.333	-	Reference
GA	0.220	0.233	0.559	1.285 (0.892-3.343)
GG	0.113	0.117	0.879	1.256(0.934-1.476)
AA	0.263	0.317	0.954	1.462(0.958-1.739)

OR: odds ratio; CI: confidence interval. Logistic regression analysis was used.

Table 6: Regression analysis for prediction of susceptibility of spontaneous recurrent abortion:


	Univariable		Multivariable	
	<i>p</i>	OR (95% CI)	<i>p</i>	OR (95% CI)
rs231775 AG+GG	0.018*	0.516(0.299-0.892)	0.001*	0.288(0.141-0.587)
rs3087243 GA+AA	0.009*	2.098(1.202-3.661)	<0.001*	3.596(1.797-7.195)

*: Significant as *P*-value <0.05; OR: odds ratio; CI: confidence interval. Logistic regression was used for analysis.

**Figure 1:** Showing normal female karyotype; 46,XX.

DISCUSSION

RPL refers to two or more consecutive clinically unsuccessful pregnancies that end before 20 full weeks of gestation, as confirmed by histology or ultrasound. It can also mean losing three or more early pregnancies in a row. Merely 2% of pregnant mothers experience two consecutive miscarriages, while as many as 50% of RPL patients lack a definitive cause (Pillarisetty and Mahdy, 2023).

**Figure 2:** LD plot in (A) control and (B) cases for studied SNPs. Squares correspond to *D'* values with numerical estimates given within the squares.

According to the current study, there is a protective effect against unexplained recurrent abortions for *CTLA-4* rs231775 AG, GG, AG+GG, and G allele genotypes, which demonstrated significantly lower frequency in cases in contrast to control groups. Our findings were supported by a meta-analysis of publications from the Chinese and PubMed databases. A total of 2607 controls and 2405 cases were included in the six case-control studies. The search criteria used to determine the abortion susceptibility associated with the *CTLA-4* +49 G/A polymorphism, there

was significantly protective association between G allele and RPL in non-Chinese population, while the reverse was found in Chinese populations (Song *et al.*, 2019).

rs231775 may reduce the risk of RPL; that is, people with the G-allele may have a lower association with RPL, or the G-allele is a protective factor for RPL risk; conversely, the A-allele is a potential risk factor, as the A-allele may rise the expression of *CTLA-4* protein, which is protective in promoting RPL and fetus toleration and this agreed with another study conducted on Han Chinese population who suggested that conduction of genetic analysis for *CTLA-4* rs231775 for RPL Patients could successfully direct efficient therapy and good parenting (Fan *et al.*, 2018); as well as other studies which were conducted on Iranian population (Rafti and Nasiri, 2016).

The National Center of Biotechnology Information's reports on the minor allele frequency (MAF) of rs3087243 for the primary global populations were examined. East Asian, 0.18; European, 0.45; African, 0.22; and South Asian, 0.60. The MAF in our analysis was 0.31 in the control group, which is slightly greater than that reported in African and East Asian populations and lower than that reported in European and Asian populations.

The polymorphism rs3087243, which is found downstream of the poly (A) termination point, may be crucial for the mRNA stability of soluble *CTLA-4*. If *CTLA-4* function is compromised, this could result in heightened T-cell activation and an increased risk of autoimmunity. (Pang *et al.*, 2022).

Our research showed that the *CTLA-4* rs3087243 genotypes and alleles were associated with the groups under investigation. Specifically, we discovered that the dominant model, GA, AA, and A alleles had significantly higher proportions in cases in contrast to the control group, which was at risk of developing recurrent abortions. In agreement with other studies in Indian Misra *et al.*, (Misra *et al.*, 2014) and Chinese population Fan *et al.* (Fan *et al.*, 2018). However, other studies reported non significant association between rs3087243 alleles and RPL in Tunisian Messaoudi *et al.*, (Messaoudi *et al.*, 2014) Japanese Hayashi *et al.* (Hayashi *et al.*, 2018); and Kazakh Svyatova *et al.* (Svyatova *et al.*, 2021), northwestern Iran Rafti and Nasiri, (Rafti and Nasiri, 2016); northern India Mojarrad *et al.*, (Mojarrad *et al.*, 2013).

In the present investigation, we noticed non significant correlation between rs231775 and rs3087243 genotypes with the number of previous abortions. Moreover, no significant association was found between *CTLA-4* rs231775-rs3087243 haplotypes with number of abortions. In consistence with others (Fan *et al.*, 2018).

Haplotype is a group of alleles that are inherited together from a single parent. Haplotypes contribute important insights into the evolutionary history of humans and could help develop more effective methods for identifying genetic variations that raise a person's risk of developing diseases (Niu, 2004).

The rs231775 and rs3087243 SNPs are spaced about 6 kilo base pairs apart. In the present study, the rs231775-rs3087243 haplotypes were calculated among each group, AG haplotype showed the highest frequency among cases (38.7%), and controls (45.9%), GG haplotype displayed the least frequency among cases (10.3%), while AA haplotype displayed the least frequency among controls (5.1%). With AG serving as the reference haplotype, GG displayed a lower frequency of cases compared to controls, offering a protective effect against the development of recurrent abortions, whereas AA displayed a higher frequency of cases in contrast to controls, indicating a susceptibility risk for recurrent abortions. GA haplotype was not associated significantly with recurrent abortion.

Our results were in agreement with others who reported GG haplotype was associated with reduced susceptibility to RPL, while AA haplotype appeared to remarkably increase the RSA risk (Fan *et al.*, 2018).

While we agreed in part with Misra and his colleagues who found that the rs231775 - rs3087243 haplotype GA carriers showed higher frequencies in RPL when compared to control groups, with risky effect to have RPL (Misra *et al.*, 2014).

Variations in sample sizes, inclusion and exclusion criteria, and ethnicities could all contribute to discrepancies in results between research. Conversely, the variability of a disease can impact the degree and kind of genetic susceptibility factors involved.

It is widely recognized that the expression of an immune-suppressing enzyme can be induced, especially in the early periods of pregnancy, by *CTLA-4* expressed on Treg cells in decidual and peripheral dendritic cells. In addition, maternal-fetal tolerance is promoted by elevated levels of the immune-suppressing enzyme (Sasaki *et al.*, 2004). Furthermore, multiple in vivo investigations found that human miscarriages were associated with down-regulated expression of Treg cells and *CTLA-4* in peripheral and decidual lymphocytes (Jin *et al.*, 2009).

One of our study's limitations is that interactions among several polymorphic loci of the same *CTLA-4* may alter the risk of RPL; these interactions should be taken into account in subsequent investigations and analyses. In addition, the sample size was insufficient to provide a higher test power. Although a number of factors influence the development

of PRL, the current study did not find any evidence of interactions between the CTLA4 gene and other genes or environmental factors. Furthermore, an assay was not conducted to measure the impact of polymorphism on the serum level of *CTLA-4*.

CONCLUSION

A recurrent *TREX1* variant rs11797 (c.531C>T) was found more frequently in lupus patients when compared to healthy controls ($p=0.01$), with minor allele frequency of 0.28 in cases and 0.16 in controls. *TREX1* positive patients had more oral ulcers, photosensitivity and seizures. However, no homozygous or heterozygous pathogenic mutations were found among this cohort of Egyptian patients. *TREX1* gene variant may play a role in the genetic susceptibility to SLE in the present cohort.

STATEMENT OF ETHICS

Our study revealed that a significant correlation was discovered between the *CTLA-4* gene rs231775 AG, GG genotypes, G allele, rs231775- rs3087243 GG haplotype with protective effect against RPL. Whereas, a significant correlation was discovered between the *CTLA-4* gene rs3087243 GA, AA genotypes a allele AA haplotype and RPL risk.

CONFLICT OF INTEREST

There are no conflicts of interest.

REFERENCES

- Abdi-Shayan S., Monfaredan A., Moradi Z., Oskoui MR., Kazemi T. (2016). Association of CD46 IVS1-1724 C> G single nucleotide polymorphism in Iranian women with unexplained recurrent spontaneous abortion (URSA). *Iranian Journal of Allergy, Asthma and Immunology*. Aug 30:303-8.
- Fan Q. e., Zhang J., Cui Y., Wang C., Xie Y., Wang Q., et al., (2018). The synergic effects of *CTLA-4*/Foxp3-related genotypes and chromosomal aberrations on the risk of recurrent spontaneous abortion among a Chinese Han population. *J. Hum Genet*, 63, 579-587.
- Hayashi Y., Nishiyama T., Nakatochi M., Suzuki S., Takahashi S. et al., (2018). Association of genetic variants of PD1 with recurrent pregnancy loss. *Reprod Med Biol*, 17, 195-202.
- Jin LP., Chen QY., Zhang T., Guo PF., L, DJ. (2009). The CD4+CD25 bright regulatory T cells and *CTLA-4* expression in peripheral and decidual lymphocytes are down-regulated in human miscarriage. *Clin Immunol*, 133, 402-10.
- MCGOWAN-JORDAN Jean, HASTINGS Ros J., MOORE Sarah, ISCN (2020): an international system for human cytogenomic nomenclature. (No Title), 2020.
- Messaoudi S., Houas I., Yaseen K., Dandana M., Mahjoub T. (2014). *CTLA-4* gene polymorphisms and risk of idiopathic recurrent pregnancy loss in a Tunisian population. *BMC Genomics*, 15, P11.
- Misra MK., Kapoor R., Pandey SK., Sharma RK., Agrawal S. (2014). Association of *CTLA-4* gene polymorphism with end-stage renal disease and renal allograft outcome. *J. Interferon Cytokine Res*, 34, 148-61.
- Mojarrad M., Hassanzadeh-Nazarabadi M., Tafazoli N. (2013). Polymorphism of genes and implantation failure. *Int J. Mol Cell Med*, 2, 1-8.
- Mumusoglu S., Beksac MS., Ekiz A., Ozdemir P., Hascelik G. (2016). Does the presence of autoantibodies without autoimmune diseases and hereditary thrombophilia have an effect on recurrent pregnancy loss?. *The Journal of Maternal-Fetal and Neonatal Medicine*. July,17;29(14):2352-7.
- Nasiri M., Rašti Z. (2016). *CTLA-4* and IL-6 gene polymorphisms: Risk factors for recurrent pregnancy loss. *Human immunology*. Dec. 1;77(12):1271-4.
- Niu T. (2004). Algorithms for inferring haplotypes. *Genet Epidemiol*, 27, 334-47.
- Pang H., Luo S., Huang G., Li X., Xie Z., Zhou Z. (2022). The Association of *CTLA-4* rs231775 and rs3087243 Polymorphisms with Latent Autoimmune Diabetes in Adults: A Meta-Analysis. *Biochem Genet*, 60, 1222-1235.
- Pillarisetty LS., Mahdy H. (2023). *Recurrent Pregnancy Loss*. StatPearls, Treasure Island (FL): StatPearls Publishing.
- Pourmasumi S., Ghasemi N., Talebi AR., Mehrabani M., Sabeti P. (2018). The effect of vitamin E and selenium on sperm chromatin quality in couples with recurrent miscarriage. *International Journal of Medical Laboratory*. Feb 10;5(1):1-0.

- Rašti Z., Nasiri M. (2016). Association of the +49 A/G Polymorphism of CTLA4 Gene with Idiopathic Recurrent Spontaneous Abortion in Women in Southwest of Iran. *J. Reprod Infertil*, 17, 151-6.
- Sasaki Y., Sakai M., Miyazaki S., Higuma S., Shiozaki A. *et al.* (2004). Decidual and peripheral blood CD4+CD25+ regulatory T cells in early pregnancy subjects and spontaneous abortion cases. *Mol Hum Reprod*, 10, 347-53.
- Song Y., Chen Y., Xu Q. (2019). Association among cytotoxic T-lymphocyte antigen 4 gene, rs231775 polymorphism, and recurrent pregnancy loss risk. *Bioscience Reports*. Feb; 39(2):BSR20181760.
- Svyatova G., Mirzakhmetova D., Berezina G., Murtazaliyeva A. (2021). Immunogenetic aspects of idiopathic recurrent miscarriage in the Kazakh population. *J. Med Life*, 14, 676-682.
- Tripathi S., Guleria I. (2015). Role of PD1/PDL1 pathway, and TH17 and treg cells in maternal tolerance to the fetus. *Biomed J.*, 38, 25-31.
- Turesheva A., Aimagambetova G., Ukybassova T., Marat A., Kanabekova P., Kaldygulova L., Amanzholkyzy A., Ryzhkova S., Nogay A., Khamidullina Z., Ilmaliyeva A. (2023). Recurrent Pregnancy Loss Etiology, Risk Factors, Diagnosis, and Management. *Fresh Look into a Full Box. Journal of Clinical Medicine*. Jun. 15;12(12):4074.
- VERMARS., BABU A. (1995). Human chromosomes: manual of basic techniques. *Chromosome Research*, 4:80-80.